arXiv:1607.08571v1 [math.SP] 28 Jul 2016

DRY TEN MARTINI PROBLEM FOR THE NON-SELF-DUAL EXTENDED HARPER'S MODEL

RUI HAN

Abstract. In this paper we prove the dry version of the Ten Martini problem: Cantor spectrum with all gaps open, for the extended Harper's model in the non self-dual region for Diophantine frequencies.

1. Introduction

The study of independent electrons on a two-dimensional lattice exposed to a perpendicular magnetic field and periodic potentials can be reduced via an appropriate choice of gauge field to the study of discrete one-dimensional quasiperiodic Jacobi matrices. The most extensively studied case is the almost Mathieu operator (AMO) acting on $l^2(\mathbb{Z})$ defined by

$$(H_{\lambda,\alpha,\theta}u)_n = u_{n+1} + u_{n-1} + 2\lambda\cos 2\pi(\theta + n\alpha)u_n.$$

This is a one-dimensional tight-binding model with anisotropic nearest neighbor couplings in general. A more general model, called the extended Harper's model (EHM), is the operator acting on $l^2(\mathbb{Z})$ defined by:

$$(H_{\lambda,\alpha,\theta}u)_n = c(\theta + n\alpha)u_{n+1} + \tilde{c}(\theta + (n-1)\alpha)u_{n-1} + 2\cos 2\pi(\theta + n\alpha)u_n.$$

where $c(\theta) = \lambda_1 e^{-2\pi i(\theta+\frac{\alpha}{2})} + \lambda_2 + \lambda_3 e^{2\pi i(\theta+\frac{\alpha}{2})}$ and $\tilde{c}(\theta) = \lambda_1 e^{2\pi i(\theta+\frac{\alpha}{2})} + \lambda_2 + \lambda_3 e^{-2\pi i(\theta+\frac{\alpha}{2})}$. It is obtained when both the nearest neighbor coupling (expressed through λ_2) and the next-nearest couplings (expressed through λ_1 and λ_3) are included. This model includes AMO as a special case (when $\lambda_1 = \lambda_3 = 0$).

For the AMO, it was proved in [5] that the spectrum is a Cantor set for any $\alpha \in \mathbb{R}\setminus\mathbb{Q}$ and $\lambda \neq 0$. This is the *Ten Martini Problem* dubbed by Barry Simon, after an offer of Mark Kac. A much more difficult problem, known as the dry version of the Ten Martini Problem, is to prove that the spectrum is not only a Cantor set, but that all gaps predicted by the Gap-Labelling theorem [10], [15] are open. The first result was obtained for Liouvillean α [12], and later it was proved for a set of (λ, α) of positive Lebesgue measure [16]. The most recent result is [6], in which they were able to deal with all Diophantine frequencies and $\lambda \neq 1$. A solution for all irrational frequencies and $\lambda \neq 1$ was also recently announced in [9].

Recently, there have been several important advances on the spectral theory of the EHM: purely point spectrum for Diophantine α and a.e.θ in the positive Lyapunov exponent region [13]; the exact formula for Lyapunov exponent for all coupling constants [14]; the spectral decomposition for a.e.α [7]. However the results that study the spectrum as a set have not been obtained for the EHM.

For EHM, depending on the values of the parameters $\lambda_1, \lambda_2, \lambda_3$, we could divide the parameter space into three regions as shown in the picture below:



$$region\ I: 0<\max(\lambda_1+\lambda_3,\lambda_2)<1,$$
$$region\ II: 0<\max(\lambda_1+\lambda_3,1)<\lambda_2,$$
$$region\ III: 0<\max(1,\lambda_2)<\lambda_1+\lambda_3.$$

According to the action of the duality transformation $\sigma:\lambda=(\lambda_1,\lambda_2,\lambda_3)\to\hat{\lambda}=(\frac{\lambda_3}{\lambda_2},\frac{1}{\lambda_2},\frac{\lambda_1}{\lambda_2})$, region I and region II are dual to each other and region III is a self-dual region. Region I is the positive Lyapunov exponent region, which is a natural extension of the segment $\{\lambda_1+\lambda_3=0,0<\lambda_2<1\}$ corresponding to the case $\lambda>1$ in the AMO. Region II is the subcritical region, which is an extension of the segment $\{\lambda_1+\lambda_3=0,1<\lambda_2\}$ corresponding to the case $\lambda<1$ in the AMO.

In this paper we prove the dry version of the Ten Martini Problem in region I and region II under the Diophantine condition.

Let p_n/q_n be the continued fraction appoximants of $\alpha\in\mathbb{R}\setminus\mathbb{Q}$. Let

$$\beta(\alpha)=\limsup_{n\to\infty}\frac{\ln q_{n+1}}{q_n}.$$

If $\beta(\alpha)=0$, we say α satisfies the Diophantine condition, denoted by $\alpha\in$ DC. It is easily seen that such α form a full measure subset of $\mathbb{T}$.

It is known that when E is in the closure of a spectral gap, the integrated density of states (IDS) $N(E)\in\alpha\mathbb{Z}+\mathbb{Z}$ (refer to (2.5) for the definition of IDS) [10], [15]. Here we prove the inverse is true.

Theorem 1.1. *If $\alpha\in$ DC and λ belongs to region I or region II, all possible spectral gaps are open.*

Remark 1.1. We note the Dry Ten Martini problem has not yet been solved for the self-dual AMO. In the self-dual region III, Cantor spectrum is known in the isotropic case (when $\lambda_1=\lambda_3$), see Fact 2.1 in [7]. In fact one could prove the operator has zero Lebesgue measure spectrum for all frequencies.

Remark 1.2. In region I and II, for Liouvillean α (where $\beta(\alpha)$ is large), it is not clear whether even the Cantor spectrum holds. The proof may require a non-trivial adjustment of the proof for AMO in [12].

We first establish almost localization (see section 3.1) in region I, then a quantitative version of Aubry duality to obtain almost reducibility (see section 3.2) in region II which enables us to deal with all energies whose rotation numbers are α-rational.

Thus the strategy follows that of [6], but we need to extend the almost localization and quantitative duality, as well as the final argument to our Jacobi setting, which is non-trivial on a technical level. At the same time unlike [6], we only deal with a short-range dual operator, leading to a significant streamlining of some arguments of [6].

We organize the paper as follows: in section 2 we present some preliminaries, in section 3 we state our main results about almost localization and almost reducibility, relying on which we provide a proof of Theorem 1.1. In section 4 and 5 we prove the main results that we present in section 3.

2. PRELIMINARIES

2.1. **Cocycles.** Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $A \in C^0(\mathbb{T}, M_2(\mathbb{C}))$ measurable with $\log \|A(x)\| \in L^1(\mathbb{T})$. The quasi-periodic *cocycle* (α, A) is the dynamical system on $\mathbb{T} \times \mathbb{C}^2$ defined by $(\alpha, A)(x, v) = (x + \alpha, A(x)v)$. The *Lyapunov exponent* is defined by

$$L(\alpha, A) = \lim_{n\to\infty} \frac{1}{n} \int_{\mathbb{T}} \log \|A_n(x)\| \mathrm{d}x = \inf_n \frac{1}{n} \int_{\mathbb{T}} \log \|A_n(x)\| \mathrm{d}x.$$

where

$$\begin{cases} A_n(x) = A(x + (n-1)\alpha) \cdots A(x) & \text{for } n \geq 0, \\ A_n(x) = A^{-1}(x + n\alpha) \cdots A^{-1}(x - \alpha) & \text{for } n < 0. \end{cases}$$

Lemma 2.1. *(e.g.*[6]*) Let (α, A) be a continous cocycle, then for any $\delta > 0$ there exists $C_\delta > 0$ such that for any $n \in \mathbb{N}$ and $\theta \in \mathbb{T}$ we have*

$$\|A_n(\theta)\| \leq C_\delta e^{(L(\alpha,A)+\delta)n}.$$

We say that (α, A) is *uniformly hyperbolic* if there exists continuous splitting $\mathbb{C}^2 = E^s(x) \bigoplus E^u(x)$, $x \in \mathbb{T}$ such that for some constant $C, \eta > 0$ and all $n \geq 0$, $\|A_n(x)v\| \leq Ce^{-\eta n}\|v\|$ for $v \in E^s(x)$ and $\|A_{-n}(x)v\| \leq Ce^{-\eta n}\|v\|$ for $v \in E^u(x)$.

Given two complex cocycles $(\alpha, A^{(1)})$ and $(\alpha, A^{(2)})$, we say they are *complex conjugate* to each other if there is $M \in C^0(\mathbb{T}, SL(2, \mathbb{C}))$ such that

$$M^{-1}(x + \alpha)A^{(1)}(x)M(x) = A^{(2)}(x).$$

We assume now that A is a real cocycle, $A \in C^0(\mathbb{T}, SL(2, \mathbb{R}))$. The notation of *real conjugacy* (between real cocycles) is the same as before, except that we look for $M \in C^0(\mathbb{T}, PSL(2, \mathbb{R}))$. A reason why we look for $M \in C^0(\mathbb{T}, PSL(2, \mathbb{R}))$ instead of $M \in C^0(\mathbb{T}, SL(2, \mathbb{R}))$ is given by the following well-known result.

Theorem 2.2. *Let (α, A) be uniformly hyperbolic, assume $\alpha \in \mathrm{DC}$ and A analytic, then there exists $M \in C^\omega(\mathbb{T}, PSL(2, \mathbb{R}))$* [1] *such that $M^{-1}(x + \alpha)A(x)M(x)$ is constant.*

We say (α, A) is (analytically) *reducible* if it is real conjugate to a constant cocycle by an analytic conjugacy.

Let

$$R_\theta = \begin{pmatrix} \cos 2\pi\theta & -\sin 2\pi\theta \\ \sin 2\pi\theta & \cos 2\pi\theta \end{pmatrix}.$$

Any $A \in C^0(\mathbb{T}, PSL(2, \mathbb{R}))$ is homotopic to $x \to R_{\frac{k}{2}x}$ for some $k \in \mathbb{Z}$ called the *degree* of A, denoted by $\deg A = k$.

[1] In general one cannot take $M \in C^\omega(\mathbb{T}, SL(2, \mathbb{R}))$.

Assume now that $A \in C^0(\mathbb{T}, SL(2,\mathbb{R}))$ is homotopic to identity. Then there exists $\phi : \mathbb{R}/\mathbb{Z} \times \mathbb{R}/\mathbb{Z} \to \mathbb{R}$ and $v : \mathbb{R}/\mathbb{Z} \times \mathbb{R}/\mathbb{Z} \to \mathbb{R}^+$ such that

$$A(x)\begin{pmatrix}\cos 2\pi y\\ \sin 2\pi y\end{pmatrix} = v(x,y)\begin{pmatrix}\cos 2\pi(y+\phi(x,y))\\ \sin 2\pi(y+\phi(x,y))\end{pmatrix}.$$

The function ϕ is called a lift of A. Let μ be any probability on $\mathbb{R}/\mathbb{Z}\times\mathbb{R}/\mathbb{Z}$ which is invariant under the continuous map $T : (x,y) \mapsto (x+\alpha, y+\phi(x,y))$, projecting over Lebesgue measure on the first coordinate. Then the number

$$\rho(\alpha, A) = \int \phi \, d\mu \bmod \mathbb{Z}$$

is independent of the choices of ϕ and μ, and is called the *fibered rotation number* of (α, A).

It can be proved directly by the definition that

$$|\rho(\alpha, A) - \theta| < C\|A - R_\theta\|_0. \tag{2.1}$$

If $(\alpha, A^{(1)})$ and $(\alpha, A^{(2)})$ are real conjugate, $M^{-1}(x+\alpha)A^{(2)}(x)M(x) = A^{(1)}(x)$, and $M : \mathbb{R}/\mathbb{Z} \to PSL(2,\mathbb{R})$ has degree k then

$$\rho(\alpha, A^{(1)}) = \rho(\alpha, A^{(2)}) - k\alpha/2. \tag{2.2}$$

For uniformly hyperbolic cocycles there is the following well-known result.

Theorem 2.3. *Let (α, A) be a uniformly hyperbolic cocycle, with $\alpha \in \mathbb{R}\backslash\mathbb{Q}$. Then $2\rho(\alpha, A) \in \alpha\mathbb{Z}+\mathbb{Z}$.*

2.2. **Extended Harper's model.** We consider the extended Harper's model $\{H_{\lambda,\theta}\}_{\theta\in\mathbb{T}}$. The formal solution to $H_{\lambda,\theta}u = Eu$ can be reconstructed via the following equation

$$\begin{pmatrix}u_{n+1}\\ u_n\end{pmatrix} = A_{\lambda,E}(\theta + n\alpha)\begin{pmatrix}u_n\\ u_{n-1}\end{pmatrix}.$$

where $A_{\lambda,E}(\theta) = \frac{1}{c(\theta)}\begin{pmatrix}E - 2\cos 2\pi\theta & -\tilde{c}(\theta-\alpha)\\ c(\theta) & 0\end{pmatrix}$. Notice that since $A_{\lambda,E}(\theta) \notin SL(2,\mathbb{R})$, we introduce the following matrix (see Lemma A.2)

$$\tilde{A}_{\lambda,E}(\theta) = \frac{1}{\sqrt{|c|(\theta)|c|(\theta-\alpha)}}\begin{pmatrix}E - 2\cos 2\pi\theta & -|c|(\theta-\alpha)\\ |c|(\theta) & 0\end{pmatrix} = Q_\lambda(\theta+\alpha)A_{\lambda,E}(\theta)Q_\lambda^{-1}(\theta),$$

where $|c|(\theta) = \sqrt{c(\theta)\tilde{c}(\theta)}$ (which is not the same as $|c(\theta)| = \sqrt{c(\theta)\overline{c(\theta)}}$ when $\theta \notin \mathbb{T}$) and $Q_\lambda(\theta)$ is analytic on $|\mathrm{Im}\theta| \le \frac{\epsilon_1}{2\pi}$.

The spectrum of $H_{\lambda,\theta}$ denoted by Σ_λ, does not depend on θ [8], and it is the set of E such that $(\alpha, \tilde{A}_{\lambda,E})$ is not uniformly hyperbolic.

The Lyapunov exponent is defined by $L_\lambda(E) = L(\alpha, A_{\lambda,E}) = L(\alpha, \tilde{A}_{\lambda,E})$.

For a matrix-valued function $M(\theta)$, let $M_\epsilon(\theta) = M(\theta + i\epsilon)$ be the phase-complexified matrix.

In [4], Avila divides all the energies in the spectrum into three catagories: super-critical, namely the energy with positive Lyapunov exponent; subcritical, namely the energy whose Lyapunov exponent of the phase-complexified cocycle is identically equal to zero in a neighborhood of $\epsilon = 0$; critical, otherwise.

The following theorem is shown in [14] (see also the appendix):

Theorem 2.4. *Extended Harper's model is super-critical in region I and sub-critical in region II. Indeed*

- *when λ belongs to region II, $L_\lambda(E) = L(\alpha, A_{\lambda,E,\epsilon}) = L(\alpha, \tilde{A}_{\lambda,E,\epsilon}) = 0$ on $|\epsilon| \le \frac{1}{2\pi}\epsilon_1(\lambda)$,*

- *when λ belongs to region II, we have $\hat{\lambda} = (\frac{\lambda_3}{\lambda_2}, \frac{1}{\lambda_2}, \frac{\lambda_1}{\lambda_2})$ belongs to region I and*

$$L_{\hat{\lambda}}(E) = \epsilon_1(\lambda), \tag{2.3}$$

where

$$\epsilon_1(\lambda) = \ln \frac{\lambda_2 + \sqrt{\lambda_2^2 - 4\lambda_1\lambda_3}}{\max(\lambda_1+\lambda_3, 1) + \sqrt{\max(\lambda_1+\lambda_3,1)^2 - 4\lambda_1\lambda_3}} > 0. \tag{2.4}$$

Fix a θ and $f \in l^2(\mathbb{Z})$. Let $\mu^f_{\lambda,\theta}$ be the *spectral measure* of $H_{\lambda,\theta}$ corresponding to f,

$$\langle (H_{\lambda,\theta} - z)^{-1} f, f\rangle = \int_{\mathbb{R}} \frac{1}{E-z} \mathrm{d}\mu^f_{\lambda,\theta}(E).$$

for z in the resolvent set $\mathbb{C} \setminus \Sigma_\lambda$.

The *integrated density of states* (IDS) is the function $N_\lambda : \mathbb{R} \to [0,1]$ defined by

$$N_\lambda(E) = \int_{\mathbb{T}} \mu^f_{\lambda,\theta}(-\infty, E] \mathrm{d}\theta, \tag{2.5}$$

where $f \in l^2(\mathbb{Z})$ is such that $\|f\|_{l^2(\mathbb{Z})} = 1$. It is a continuous non-decreasing surjective funtion.

Notice that $\tilde{A}_{\lambda,E}(\theta) \in SL(2,\mathbb{R})$ is homotopic to identity in $C^0(\mathbb{T}, SL(2,\mathbb{R}))$, in fact just consider

$$H_t(\lambda, E, \theta) = \frac{1}{\sqrt{|c|(\theta)|c|(\theta - t\alpha)}} \begin{pmatrix} t(E - v(\theta)) & -|c|(\theta - t\alpha) \\ |c|(\theta) & 0 \end{pmatrix}.$$

which establishes a homotopy of $\tilde{A}_{\lambda,E}(\theta)$ to $R_{\frac{1}{4}}$ and hence to the identity. Therefore we can define the rotation number $\rho(\alpha, \tilde{A}_{\lambda,E})$. Let $\rho_\lambda(E) = \rho(\alpha, \tilde{A}_{\lambda,E})$. Notice that $\rho_\lambda(E)$ is associated to the operator

$$(\tilde{H}_{\lambda,\theta} u)_n = |c|(\theta + n\alpha)u_{n+1} + |c|(\theta + (n-1)\alpha)u_{n-1} + 2\cos 2\pi(\theta + n\alpha)u_n.$$

It is easily seen that for each θ, $\tilde{H}_{\lambda,\theta}$ and $H_{\lambda,\theta}$ differ by a unitary operator, thus they share the same spectrum and integrated density of states, $\tilde{N}_\lambda(E) = N_\lambda(E)$. The relation between the integrated density of states and rotation number of $\tilde{H}_{\lambda,\theta}$ yields the following

$$N_\lambda(E) = \tilde{N}_\lambda(E) = 1 - 2\rho_\lambda(E). \tag{2.6}$$

2.3. **The dual model.** It turns out the spectrum Σ_λ of $H_{\lambda,\theta}$ is related to the spectrum $\Sigma_{\hat{\lambda}}$ of $H_{\hat{\lambda},\theta}$ in the following way

$$\Sigma_\lambda = \lambda_2 \Sigma_{\hat{\lambda}}$$

by Aubry duality. This map $\sigma : \lambda \to \hat{\lambda}$ establishes the duality between region I and region II. The IDS $N_\lambda(E)$ of $H_{\lambda,\theta}$ coincide with the IDS $N_{\hat{\lambda}}(E/\lambda_2)$ of $H_{\hat{\lambda},\theta}$. Since $\Sigma_\lambda = \lambda_2\Sigma_{\hat{\lambda}}$, we have the following

Theorem 2.5. [11], [17] *For any λ, θ, there exists a dense set of $E \in \Sigma_\lambda$ such that there exists a non-zero solution of $H_{\hat{\lambda},\theta} u = \frac{E}{\lambda_2} u$ with $|u_k| \le 1 + |k|$.*

2.4. **Bounded eigenfunction for every energy.** The next result from [6] allows us to pass from a statement of every θ to every E.

Theorem 2.6. [6] *If $E \in \Sigma_\lambda$ then there exists $\theta(E) \in \mathbb{T}$ and a bounded solution of $H_{\hat{\lambda},\alpha,\theta} u = \frac{E}{\lambda_2} u$ with $u_0 = 1$ and $|u_k| \le 1$.*

2.5. **Localization and reducibility.**

Theorem 2.7. *Given α irrational, $\theta \in \mathbb{R}$ and λ in region II, fix $E \in \Sigma_\lambda$, and suppose $H_{\hat{\lambda},\theta} u = \frac{E}{\lambda_2} u$ has a non-zero exponentially decaying eigenfunction $u = \{u_k\}_{k\in\mathbb{Z}}$, $|u_k| \le e^{-c|k|}$ for k large enough. Then the following hold:*

- *(A) If $2\theta \notin \alpha\mathbb{Z} + \mathbb{Z}$, then there exists $M : \mathbb{R}/\mathbb{Z} \to SL(2,\mathbb{R})$ analytic, such that*
$$M^{-1}(x+\alpha)\tilde{A}_{\lambda,E}(x)M(x) = R_{\pm\theta}.$$
In this case $\rho(\alpha, \tilde{A}_{\lambda,E}) = \pm\theta + \frac{m}{2}\alpha \bmod \mathbb{Z}$, where $m = \deg M$ (here since $M \in SL(2,\mathbb{R})$, we have that m is an even number) and $2\rho(\alpha, \tilde{A}_{\lambda,E}) \notin \alpha\mathbb{Z} + \mathbb{Z}$.
- *(B) If $2\theta \in \alpha\mathbb{Z} + \mathbb{Z}$ and $\alpha \in$ DC, then there exists $M : \mathbb{R}/\mathbb{Z} \to PSL(2,\mathbb{R})$ analytic, such that*
$$M^{-1}(x+\alpha)\tilde{A}_{\lambda,E}(x)M(x) = \begin{pmatrix} \pm 1 & a \\ 0 & \pm 1 \end{pmatrix}$$
with $a \neq 0$. In this case $\rho(\alpha, \tilde{A}_{\lambda,E}) = \frac{m}{2}\alpha \bmod \mathbb{Z}$, where $m = \deg M$, i.e. $2\rho(\alpha, \tilde{A}_{\lambda,E}) \in \alpha\mathbb{Z} + \mathbb{Z}$.

Proof. Let $u(x) = \sum_{k\in\mathbb{Z}} \hat{u}_k e^{2\pi i k x}$, $U(x) = \begin{pmatrix} e^{2\pi i\theta} u(x) \\ u(x-\alpha) \end{pmatrix}$. Then

$$A_{\lambda,E}(x)U(x) = e^{2\pi i\theta} U(x+\alpha),$$

$$\tilde{A}_{\lambda,E}(x)\tilde{U}(x) = e^{2\pi i\theta}\tilde{U}(x+\alpha).$$

Notice $\tilde{U}(x) = Q_\lambda(x)U(x)$ is analytic in $|\mathrm{Im} x| < \frac{\tilde{c}}{2\pi}$, where $\tilde{c} = \min(\epsilon_1, c)$, ϵ_1 as in 2.4 and Q_λ as in A.2. Define $\overline{\tilde{U}(x)}$ to be the complex conjugate of $\tilde{U}(x)$ on $\mathbb{T}$ and its analytic extension to $|\mathrm{Im} x| < \frac{\tilde{c}}{2\pi}$. Let $M(x)$ be the matrix with columns $\tilde{U}(x)$ and $\overline{\tilde{U}(x)}$. Then,

$$\tilde{A}_{\lambda,E}(x)M(x) = M(x+\alpha)\begin{pmatrix} e^{2\pi i\theta} & 0 \\ 0 & e^{-2\pi i\theta} \end{pmatrix} \quad \text{on } \mathbb{T}.$$

Then since $\det M(x+\alpha) = \det M(x)$, we know $\det M(x)$ is a constant on $\mathbb{T}$.

Case 1. If $\det M(x) \neq 0$, then let $M(x) = \tilde{M}(x)\begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix}$.

$$\tilde{M}^{-1}(x+\alpha)\tilde{A}_{\lambda,E}(x)\tilde{M}(x) = R_\theta = \begin{pmatrix} \cos 2\pi\theta & -\sin 2\pi\theta \\ \sin 2\pi\theta & \cos 2\pi\theta \end{pmatrix}.$$

Case 2. If $\det M(x) = 0$, then if we denote $\tilde{U}(x) = \begin{pmatrix} u_1(x) \\ u_2(x) \end{pmatrix}$, then $\det M(x) = 0$ means there exists $\eta(x)$ such that $u_1(x) = \eta(x)\overline{u_1(x)}$ and $u_2(x) = \eta(x)\overline{u_2(x)}$. This implies that $\eta(x) \in \mathbb{C}^\omega(\mathbb{T}, \mathbb{C})$, and $|\eta(x)| = 1$ on $\mathbb{T}$. Therefore there exists $\phi(x) \in \mathbb{C}^\omega(\mathbb{R}/2\mathbb{Z}, \mathbb{C})$ such that $\phi^2(x) = \eta(x)$ and $|\phi(x)| = 1$. It is easy to see $\overline{\phi(x)}u_1(x) = \phi(x)\overline{u_1(x)}$ and $\overline{\phi(x)}u_2(x) = \phi(x)\overline{u_2(x)}$. Then we define $W(x) = \begin{pmatrix} \overline{\phi(x)}u_1(x) \\ \overline{\phi(x)}u_2(x) \end{pmatrix}$, it is a real vector on $\mathbb{R}/2\mathbb{Z}$ with $W(x+1) = \pm W(x)$, and $\tilde{U}(x) = \phi(x)W(x)$. Now let us define $\tilde{M}(x)$ to be the matrix with columns $W(x)$ and $\frac{1}{\|W(x)\|^{-2}} R_{\frac{1}{4}} W(x)$, then $\det \tilde{M}(x) = 1$ and $\tilde{M}(x) \in PSL(2,\mathbb{R})$. Since

$$\tilde{A}_{\lambda,E}(x)W(x) = \frac{e^{2\pi i\theta}\phi(x+\alpha)}{\phi(x)} W(x+\alpha).$$

We have

$$\tilde{A}_{\lambda,E}(x)\tilde{M}(x) = \tilde{M}(x+\alpha)\begin{pmatrix} d(x) & \tau(x) \\ 0 & d(x)^{-1}\end{pmatrix}$$

where $d(x) = \frac{e^{2\pi i\theta}\phi(x+\alpha)}{\phi(x)}$, $|d(x)| = 1$ and $d(x)$ being real number, therefore $d(x) = \pm 1$. Also $\tau(x) \in \mathbb{C}^\omega(\mathbb{R}/2\mathbb{Z}, \mathbb{C})$. But in fact $\tilde{M}^{-1}(x+\alpha)\tilde{A}_{\lambda,E}(x)\tilde{M}(x)$ is well-defined on $\mathbb{T}$. Therefore $\tau(x) \in \mathbb{C}^\omega(\mathbb{T}, \mathbb{C})$. Now since we assumed $\alpha \in$ DC, we can further reduce $\tau(x)$ to the constant $\tau = \int_{\mathbb{T}} \tau(x)\mathrm{d}x$. In fact there exists $\psi(x) \in \mathbb{C}^\omega(\mathbb{T}, \mathbb{C})$ such that $-\psi(x+\alpha) + \psi(x) + \tau(x) = \int_{\mathbb{T}} \tau(x)\mathrm{d}x$. This implies

$$\begin{pmatrix} 1 & -\psi(x+\alpha) \\ 0 & 1\end{pmatrix}\tilde{M}^{-1}(x+\alpha)\tilde{A}_{\lambda,E}(x)\tilde{M}(x)\begin{pmatrix} 1 & \psi(x) \\ 0 & 1\end{pmatrix} = \begin{pmatrix} \pm 1 & \tau \\ 0 & \pm 1\end{pmatrix}.$$

In fact if $\det M(x) = 0$, then $\frac{e^{2\pi i\theta}\phi(x+\alpha)}{\phi(x)} = \pm 1$, which implies that $2\theta \in \alpha\mathbb{Z} + \mathbb{Z}$. Therefore if $2\theta \notin \alpha\mathbb{Z} + \mathbb{Z}$, we must be in case (A). If on the other hand, $2\theta \in \alpha\mathbb{Z} + \mathbb{Z}$, $2\theta = k\alpha + n$, suppose $\tilde{M}^{-1}(x+\alpha)\tilde{A}_{\lambda,E}(x)\tilde{M}(x) = R_\theta$, then $R_{-\frac{k}{2}(x+\alpha)}\tilde{M}^{-1}(x+\alpha)\tilde{A}_{\lambda,E}(x)\tilde{M}(x)R_{\frac{k}{2}x} = R_{\frac{n}{2}} = \pm I$ leading to a contradiction. Therefore if $2\theta \in \alpha\mathbb{Z} + \mathbb{Z}$, we must be in case (B). □

2.6. **Continued fractions.** Let $\{q_n\}$ be the denominators of the continued fraction approximants of α. We recall the following properties:

$$\|q_n\alpha\|_{\mathbb{R}/\mathbb{Z}} = \inf_{1\le|k|\le q_{n+1}-1}\|k\alpha\|_{\mathbb{R}/\mathbb{Z}},$$

$$\frac{1}{2q_{n+1}} \le \|q_n\alpha\|_{\mathbb{R}/\mathbb{Z}} \le \frac{1}{q_{n+1}}.$$

Recall that the Diophantine condition of α is $\beta(\alpha) = \limsup_{n\to\infty}\frac{\ln q_{n+1}}{q_n} = 0$. Thus for any $\xi > 0$, there exists $C_\xi > 0$ such that

$$\|k\alpha\|_{\mathbb{R}/\mathbb{Z}} \ge C_\xi e^{-\xi|k|} \quad \text{for any } k \ne 0. \tag{2.7}$$

Lemma 2.8. [5] *Let $\alpha \in \mathbb{R}\backslash\mathbb{Q}$, $x \in \mathbb{R}$ and $0 \le l_0 \le q_n - 1$ be such that $|\sin\pi(x+l_0\alpha)| = \inf_{0\le l\le q_n-1}|\sin\pi(x+l\alpha)|$, then for some absolute constant $C_1 > 0$,*

$$-C_1\ln q_n \le \sum_{0\le l\le q_n-1, l\ne l_0}\ln|\sin\pi(x+l\alpha)| + (q_n-1)\ln 2 \le C_1\ln q_n$$

Lemma 2.9. [6] *Let $1 \le r \le [q_{n+1}/q_n]$. If $p(x)$ has essential degree at most $k = rq_n - 1$ and $x_0 \in \mathbb{R}/\mathbb{Z}$, then for some absolute constant C_2,*

$$\|p(x)\|_0 \le C_2 q_{n+1}^{C_2 r}\sup_{0\le j\le k}|p(x_0+j\alpha)|.$$

3. Main estimates and proof of Theorem 1.1

3.1. **Almost localization for every θ.**

Definition 3.1. Let $\alpha \in \mathbb{R}\setminus\mathbb{Q}$, $\theta \in \mathbb{R}$, $\epsilon_0 > 0$. We say that k is an ϵ_0-resonance of θ if $\|2\theta - k\alpha\| \le e^{-\epsilon_0|k|}$ and $\|2\theta - k\alpha\| = \min_{|l|\le|k|}\|2\theta - l\alpha\|$.

Definition 3.2. Let $0 = |n_0| < |n_1| < ...$ be the ϵ_0-resonances of θ. If this sequence is infinite, we say θ is ϵ_0-resonant, otherwise we say it is ϵ_0-non-resonant.

Definition 3.3. We say the extended Harper's model $\{H_{\lambda,\alpha,\theta}\}_\theta$ exhibits almost localization if there exists $C_0, C_3, \epsilon_0, \tilde{\epsilon}_0 > 0$, such that for every solution ϕ to $H_{\lambda,\alpha,\theta}\phi = E\phi$ satisfying $\phi(0) = 1$ and $|\phi(m)| \leq 1 + |m|$, and for every $C_0(1 + |n_j|) < |k| < C_0^{-1}|n_{j+1}|$, we have $|\phi(k)| \leq C_3 e^{-\tilde{\epsilon}_0|k|}$ (where n_j are the ϵ_0-resonances of θ).

Theorem 3.1. *If λ belongs to region II, $\{H_{\hat{\lambda},\alpha,\theta}\}_\theta$ is almost localized for every $\alpha \in$ DC.*

Remark 3.1. It is clear from Theorem 3.1 that almost localization implies localization for non-resonant θ.

We will actually prove the following explicit lemma:

Lemma 3.2. *Let λ be in region II. Let C_4 be the absolute constant in Lemma* 4.3*, $\epsilon_1 = \epsilon_1(\lambda)$ be as in (*2.4*), then for any $0 < \epsilon_0 < \frac{\epsilon_1}{100C_4}$, there exists constant $C_3 > 0$, which depends on λ, α and ϵ_0, so that for every solution u of $H_{\hat{\lambda},\alpha,\theta}u = Eu$ satisfying $u(0) = 1$ and $|u_k| \leq 1 + |k|$, if $3(|n_j| + 1) < |k| < \frac{1}{3}|n_{j+1}|$, then $|u_k| \leq C_3 e^{-\frac{\epsilon_1}{5}|k|}$, where $\{n_j\}$ are the ϵ_0-resonances of θ.*

The proof of Lemma 3.2 (and thus of Theorem 3.1) is given in Section 4.

3.2. **Almost reducibility.**

Let λ be in region II. For every $E \in \Sigma_\lambda$, let $\theta(E) \in \mathbb{T}$ be given in Theorem 2.6. Let $0 < \epsilon_0 < \frac{\epsilon_1}{100C_4}$ and $\{n_j\}$ be the set of ϵ_0- resonances of $\theta(E)$. Then for some positive constants N_0, C and c, independent of E and θ, we have the following theorem:

Theorem 3.3. *For any fixed j, with $N_0 < n = |n_j| + 1 < \infty$, let $N = |n_{j+1}|$, $L^{-1} = \|2\theta - n_j\alpha\|$. Then there exists $W : \mathbb{T} \to SL(2, \mathbb{R})$ analytic such that $|\deg W| \leq Cn$, $\|W\|_0 \leq CL^C$ and $\|W^{-1}(x + \alpha)\tilde{A}_{\lambda,E}(x)W(x) - R_{\mp\theta}\| \leq Ce^{-cN}$.*

Remark 3.2. Notice that this theorem requires $n > N_0$, which is not always ensured when $\theta(E)$ is non-resonant, however in that case we have localization for $H_{\hat{\lambda},\alpha,\theta}$ instead of almost localization. We will prove Theorem 3.3 in Section 5.

3.3. **Spectral consequences of Almost reducibility.**

Let $\epsilon_1 = \epsilon_1(\lambda)$ and C_4 be as in Lemma 3.2.

Theorem 3.4. *Assume $\alpha \in$ DC. For λ in region II, fix $E \in \Sigma_\lambda$. Assume $\theta(E) \in \mathbb{T}$ is such that $H_{\hat{\lambda},\alpha,\theta}u = \frac{E}{\lambda_2}u$ has solution satisfying $u_0 = 1$ and $|u_k| \leq 1$. Let C be the constant in Theorem* 3.3*. Then $\theta(E)$ and $\rho(\alpha, \tilde{A}_{\lambda,E})$ have the following relation:*

- *(A) If θ is ϵ_0-non-resonant for some $\frac{\epsilon_1}{100C_4} > \epsilon_0 > 0$, then $2\theta \in \mathbb{Z}\alpha + \mathbb{Z}$ if and only if $2\rho(\alpha, \tilde{A}_{\lambda,E}) \in \mathbb{Z}\alpha + \mathbb{Z}$.*
- *(B) If θ is ϵ_0-resonant for some $\frac{\epsilon_1}{100C_4} > \epsilon_0 > 0$, then $\rho(\alpha, \tilde{A}_{\lambda,E})$ is $\frac{\epsilon_0}{C+2}$-resonant.*

Proof.

(A): When θ is ϵ_0-non-resonant for some $\frac{\epsilon_1}{100C_4} > \epsilon_0 > 0$, Theorem 3.1 implies $H_{\hat{\lambda},\alpha,\theta}$ has exponentially decaying eigenfunction. Then applying Theorem 2.7 we get $2\theta \in \mathbb{Z}\alpha + \mathbb{Z}$ if and only if $2\rho(\alpha, \tilde{A}_{\lambda,E}) \in \mathbb{Z}\alpha + \mathbb{Z}$.

(B): Assume θ is ϵ_0-resonant for some $\frac{\epsilon_1}{100C_4} > \epsilon_0 > 0$. Fix any $\xi < \frac{\epsilon_0}{2C+2}$, then there exists $C_\xi > 0$ such that for any $k \neq 0$ we have $\|k\alpha\| \geq C_\xi e^{-\xi|k|}$. Now take an ϵ_0-resonance n_j of θ such that $n = |n_j| > \max\left(\frac{-\ln C_\xi/2}{\epsilon_0 - (2C+2)\xi}, N_0\right)$. Then there exists $|m| \leq Cn$ such that $2\rho(\alpha, \tilde{A}_{\lambda,E}) - m\alpha = -2\theta$. Then

$$\|2\rho(\alpha, \tilde{A}_{\lambda,E}) - (m - n_j)\alpha\| = \|2\theta - n_j\alpha\| < e^{-\epsilon_0 n} \leq e^{-\frac{\epsilon_0}{C+2}|m-n_j|}.$$

Take any $|l| \leq |m-n_j|$, $l \neq m-n_j$. Then

$$\|(l-(m-n_j))\alpha\| \geq C_\xi e^{-2\xi|m-n_j|} > 2e^{-\epsilon_0 n} > 2\|2\rho(\alpha,\tilde{A}_E)-(m-l_0)\alpha\|.$$

Thus $\|2\rho(\alpha,\tilde{A}_E)-l\alpha\| > \|2\rho(\alpha,\tilde{A}_E)-(m-n_j)\alpha\|$ for any $|l| \leq |m-n_j|$, $l \neq m-n_j$. This by definition means $\rho(\alpha,\tilde{A}_{\lambda,E})$ is $\frac{\epsilon_0}{C+2}$-resonant. □

Now based on Theorem 3.4, we can complete the proof of the dry version of Ten Martini Problem for extended Harper's model in regions I and II.

Proof of Theorem 1.1

It is enough to consider λ in region II. Let $E \in \Sigma_\lambda$ be such that $N_\lambda(E) \in \mathbb{Z}\alpha+\mathbb{Z}$. We are going to show E belongs to the boundary of a component of $\mathbb{R}\setminus\Sigma_\lambda$. Now by (2.6) we have $2\rho(\alpha,\tilde{A}_{\lambda,E}) \in \alpha\mathbb{Z}+\mathbb{Z}$, thus by Theorem 3.4, $2\theta(E) \in \alpha\mathbb{Z}+\mathbb{Z}$. By Theorem 2.7, this means there exist $M(x) \in C_h^\omega(\mathbb{T}, PSL(2,\mathbb{R}))$ such that $M^{-1}(x+\alpha)\tilde{A}_{\lambda,E}(x)M(x) = \begin{pmatrix}\pm 1 & a\\ 0 & \pm 1\end{pmatrix}$. Without loss of generality, we assume $M^{-1}(x+\alpha)\tilde{A}_{\lambda,E}(x)M(x) = \begin{pmatrix}1 & a\\ 0 & 1\end{pmatrix}$. Let $\tilde{M}(x) = \frac{M(x)}{\sqrt{|c|(x-\alpha)}}$, then

$$\tilde{M}^{-1}(x+\alpha)\begin{pmatrix}\frac{E-v(x)}{|c|(x)} & -\frac{|c|(x-\alpha)}{|c|(x)}\\ 1 & 0\end{pmatrix}\tilde{M}(x) = \begin{pmatrix}1 & a\\ 0 & 1\end{pmatrix}.$$

Now let $\tilde{M}(x) = \begin{pmatrix}M_{11}(x) & M_{12}(x)\\ M_{21}(x) & M_{22}(x)\end{pmatrix}$. Then $M_{21}(x) = M_{11}(x-\alpha)$ and $M_{22}(x) = M_{12}(x-\alpha) - aM_{11}(x-\alpha)$ amd

$$\begin{aligned}&\tilde{M}^{-1}(x+\alpha)\begin{pmatrix}\frac{E+\epsilon-v(x)}{|c|(x)} & -\frac{|c|(x-\alpha)}{|c|(x)}\\ 1 & 0\end{pmatrix}\tilde{M}(x)\\ =&\begin{pmatrix}1 & a\\ 0 & 1\end{pmatrix} + \epsilon\begin{pmatrix}M_{11}(x)M_{12}(x)-aM_{11}^2(x) & M_{12}^2(x)-aM_{11}(x)M_{12}(x)\\ -M_{11}^2(x) & -M_{11}(x)M_{12}(x)\end{pmatrix}.\\ \triangleq& M_0+\epsilon M_1(x).\end{aligned}$$

Now we look for $Z_\epsilon(x)$ of the form $e^{\epsilon Y(x)}$ such that

$$Z_\epsilon^{-1}(x+\alpha)(M_0+\epsilon M_1(x))Z_\epsilon(x) = M_0+\epsilon[M_1]+O(\epsilon^2).$$

We then just need to solve the equation:

$$(I-\epsilon Y(x+\alpha)+O(\epsilon^2))(M_0+\epsilon M_1(x))(I+\epsilon Y(x)+O(\epsilon^2)) = M_0+\epsilon[M_1]+O(\epsilon^2).$$

It is sufficient to solve the coholomogical equation:

$$Y(x+\alpha)M_0 - M_0Y(x) = M_1(x)-[M_1],$$

which is guaranteed by the Diophantine condition on α. Thus

$$\begin{aligned}&(M(x+\alpha)Z_\epsilon(x+\alpha))^{-1}\tilde{A}_{\lambda,E}(x)(M(x)Z_\epsilon(x))\\ =&\begin{pmatrix}1+\epsilon[M_{11}M_{12}]-a\epsilon[M_{11}^2] & a+\epsilon[M_{12}^2]-a\epsilon[M_{11}M_{12}]\\ -\epsilon[M_{11}^2] & 1-\epsilon[M_{11}M_{12}]\end{pmatrix}+O(\epsilon^2)\\ \triangleq& M_\epsilon+O(\epsilon^2).\end{aligned}$$

Notice that $\tilde{A}_{\lambda,E}$ is uniformly hyperbolic iff $\mathrm{Trace}(M_\epsilon) > 2$ which is fulfilled when $-a\epsilon[M_{11}^2] > 0$. Thus for ϵ small, satisfying $-a\epsilon[M_{11}^2] > 0$, $E+\epsilon \notin \Sigma_\lambda$, which means this spectral gap is open. □

4. Almost localization in region I

In this section we will prove Lemma 3.2. For fixed λ in region II and E, let $D_{\hat{\lambda},E}(\theta) = c_{\hat{\lambda}}(\theta)A_{\hat{\lambda},E}(\theta)$, where $c_{\hat{\lambda}}(\theta) = \frac{\lambda_3}{\lambda_2}e^{-2\pi i(\theta+\frac{\alpha}{2})} + \frac{1}{\lambda_2} + \frac{\lambda_1}{\lambda_2}e^{2\pi i(\theta+\frac{\alpha}{2})}$. Regarding the Lyapunov exponent, we recall the following result in [14],

$$L(\alpha, A_{\hat{\lambda},E}) = L(\alpha, D_{\hat{\lambda},E}) - \int_{\mathbb{T}} \ln|c_{\hat{\lambda}}(\theta)|\mathrm{d}\theta \triangleq \tilde{L} - \int \ln|c_{\hat{\lambda}}| > 0,$$

where $\tilde{L} = \ln\frac{\lambda_2+\sqrt{\lambda_2^2-4\lambda_1\lambda_3}}{2\lambda_2}$ and $\int \ln|c_{\hat{\lambda}}| = \ln\frac{\max(\lambda_1+\lambda_3,1)+\sqrt{\max(\lambda_1+\lambda_3,1)^2-4\lambda_1\lambda_3}}{2\lambda_2}$.

Proof of of Lemma 3.2

Suppose u is a solution satisfying the condition of Lemma 3.2. For an interval $I = [x_1, x_2]$, let Γ_I be the coupling operator between I and $\mathbb{Z}\setminus I$:

$$\Gamma_I(i,j) = \begin{cases} \tilde{c}(\theta+(x_1-1)\alpha), & (i,j) = (x_1, x_1-1) \\ c(\theta+(x_1-1)\alpha), & (i,j) = (x_1-1, x_1) \\ \tilde{c}(\theta+x_2\alpha), & (i,j) = (x_2+1, x_2) \\ c(\theta+x_2\alpha), & (i,j) = (x_2, x_2+1) \\ 0 & \text{otherwise.} \end{cases}$$

Let $H_I = R_I H_{\hat{\lambda},\theta} R_I^*$ be the restricted operator of $H_{\hat{\lambda},\theta}$ to I. Then for $x \in I$, we have $(H_I + \Gamma_I - E)u(x) = 0$. Thus $u(x) = G_I\Gamma_I u(x)$, where $G_I = (E-H_I)^{-1}$. By matrix multiplication:

$$\begin{aligned} u(x) &= \sum_{y\in I,(y,z)\in\Gamma_I} G_I(x,y)\Gamma_I(y,z)u(z) \\ &= \tilde{c}(\theta+(x_1-1)\alpha)G_I(x,x_1)u(x_1-1) + c(\theta+x_2\alpha)G_I(x,x_2)u(x_2+1). \end{aligned}$$

Let us denote $P_k(\theta) = \det(E - H_{[0,k-1]}(\theta))$. Then the $k-$step matrix $D_{\hat{\lambda},E,k}(\theta)$ satisfies:

$$D_{\hat{\lambda},E,k}(\theta) = \begin{pmatrix} P_k(\theta) & -\tilde{c}(\theta-\alpha)P_{k-1}(\theta+\alpha) \\ c(\theta+(k-1)\alpha)P_{k-1}(\theta) & -\tilde{c}(\theta-\alpha)c(\theta+(k-1)\alpha)P_{k-2}(\theta+\alpha) \end{pmatrix}.$$

This relation between $P_k(\theta)$ and $D_{\hat{\lambda},E,k}(\theta)$ gives a general upper bound of $P_k(\theta)$ in terms of $\tilde{L}$. Indeed by Lemma 2.1, for any $\epsilon > 0$ there exists $C(\epsilon) > 0$ so that

$$|P_n(\theta)| \le C(\epsilon)e^{(\tilde{L}+\epsilon)n} \quad \text{for any } n \in \mathbb{N}.$$

By Cramer's rule:

$$|G_I(x_1,y)| = \prod_{j=x_1}^{y-1}|c(\theta+j\alpha)|\left|\frac{\det(E-H_{[y+1,x_2]}(\theta))}{\det(E-H_I(\theta))}\right| = \prod_{j=x_1}^{y-1}|c(\theta+j\alpha)|\left|\frac{P_{x_2-y}(\theta+(y+1)\alpha)}{P_k(\theta+x_1\alpha)}\right|,$$

$$|G_I(y,x_2)| = \prod_{j=y+1}^{x_2}|c(\theta+j\alpha)|\left|\frac{\det(E-H_{[x_1,y-1]}(\theta))}{\det(E-H_I(\theta))}\right| = \prod_{j=y+1}^{x_2}|c(\theta+j\alpha)|\left|\frac{P_{y-x_1}(\theta+x_1\alpha)}{P_k(\theta+x_1\alpha)}\right|.$$

Notice that $P_k(\theta)$ is an even function about $\theta+\frac{k-1}{2}\alpha$, it can be written as a polynomial of degree k in $\cos 2\pi(\theta+\frac{k-1}{2}\alpha)$. Let $P_k(\theta) = Q_k(\cos 2\pi(\theta+\frac{k-1}{2}\alpha))$. Let $M_{k,r} = \{\theta\in\mathbb{T},\ |Q_k(\cos 2\pi\theta)| \le e^{(k+1)r}\}$.

Definition 4.1. Fix $m > 0$. A point $y \in \mathbb{Z}$ is called $(k,m)-$regular if there exists an interval $[x_1,x_2]$ containing y, where $x_2 = x_1+k-1$ such that

$$|G_I(y,x_i)| \le e^{-m|y-x_i|} \text{ and } \operatorname{dist}(y,x_i) \ge \frac{1}{3}k \text{ for } i = 1,2,$$

otherwise y is called $(k,m)-$singular.

Lemma 4.1. *Suppose $y\in\mathbb{Z}$ is $(k,\tilde{L}-\int\ln|c_{\hat{\lambda}}|-\rho)-$singular. Then for any $\epsilon>0$ and any $x\in\mathbb{Z}$ satisfying $y-\frac{2}{3}k\le x\le y-\frac{1}{3}k$, we have $\theta+(x+\frac{1}{2}(k-1))\alpha$ belongs to $M_{k,\tilde{L}-\frac{1}{3}\rho+\epsilon}$ for $k>k(\lambda,\epsilon,\rho)$.*

Proof. Suppose there exists $\epsilon>0$ and x_1: $y-(1-\delta)k\le x_1\le y-\delta k$, such that $\theta+(x_1+\frac{1}{2}(k-1))\alpha$ does not belong to $M_{k,\tilde{L}-\frac{1}{3}\rho+\epsilon}$, that is $|P_k(\theta+x_1\alpha)|>e^{(k+1)(\tilde{L}-\rho\delta+\epsilon)}$,

$$\begin{aligned}|G_I(x_1,y)|&\le\prod_{j=x_1}^{y-1}|c_{\hat{\lambda}}(\theta+j\alpha)|e^{(k-|x_1-y|)(\tilde{L}+\epsilon)}e^{-(k+1)(\tilde{L}-\frac{1}{3}\rho+\epsilon)}\\&<e^{-(\tilde{L}-\int\ln|c_{\hat{\lambda}}|-\rho)|y-x_1|}\ \text{ for } k>k(\lambda,\epsilon,\rho).\end{aligned}$$

Similarly

$$|G_I(x_2,y)|\le e^{-(\tilde{L}-\int\ln|c_{\hat{\lambda}}|-\rho)|y-x_2|}.$$

□

Definition 4.2. We say that the set $\{\theta_1,...,\theta_{k+1}\}$ is $\gamma-$uniform if

$$\max_{x\in[-1,1]}\max_{i=1,...,k+1}\prod_{j=1,j\neq i}^{k+1}\frac{|x-\cos2\pi\theta_j|}{|\cos2\pi\theta_i-\cos2\pi\theta_j|}<e^{k\gamma}$$

Lemma 4.2. *Let $\gamma_1<\gamma$. If $\theta_1,...,\theta_{k+1}\in M_{k,\tilde{L}-\gamma}$, then $\{\theta_1,...,\theta_{k+1}\}$ is not γ_1-uniform for $k>k(\gamma,\gamma_1)$.*

Proof. Otherwise, using Lagrange interpolation form we can get $|Q_k(x)|<e^{k\tilde{L}}$ for all $x\in[-1,1]$. This implies $|P_k(x)|<e^{k\tilde{L}}$ for all x. But by Herman's subharmonic function argument, $\int_{\mathbb{R}/\mathbb{Z}}\ln|P_k(x)|\mathrm{d}x\ge k\tilde{L}$. This is impossible. □

Now take ξ and ϵ_0 such that $0<1000\xi<\epsilon_0$. Then for $|n_{j+1}|>N(\xi)$ we have

$$2e^{-4\xi|n_{j+1}|}\le C_\xi e^{-2\xi|n_{j+1}|}\le\|(n_{j+1}-n_j)\alpha\|=\|n_{j+1}\alpha-2\theta+2\theta-n_j\alpha\|\le2\|2\theta-n_j\alpha\|\le2e^{-\epsilon_0|n_j|},$$

which yields that

$$|n_{j+1}|>\frac{\epsilon_0}{4\xi}|n_j|>250|n_j|. \tag{4.1}$$

Without loss of generality, assume $3(|n_j|+1)<y<\frac{|n_{j+1}|}{3}$ and $y>N(\xi)$. Select n such that $q_n\le\frac{y}{8}<q_{n+1}$ and let s be the largest positive integer satisfying $sq_n\le\frac{y}{8}$. Set $I_1,I_2\subset\mathbb{Z}$ as follows

$$\begin{aligned}&I_1=[1-2sq_n,0]\ \textit{and}\ I_2=[y-2sq_n+1,y+2sq_n],\ \text{if } n_j<0\\&I_1=[0,2sq_n-1]\ \textit{and}\ I_2=[y-2sq_n+1,y+2sq_n],\ \text{if } n_j\ge0\end{aligned}$$

Lemma 4.3. *Let $\theta_j=\theta+j\alpha$, then set $\{\theta_j\}_{j\in I_1\cup I_2}$ is $C_4\epsilon_0+C_4\xi-$uniform for some absolute constant C_4 and $y>y(\alpha,\epsilon_0,\xi)$.*

Proof. Without loss of generality, we assume $n_j > 0$. Take $x = \cos 2\pi a$. Now it suffices to estimate

$$\sum_{j\in I_1\cup I_2,\ j\neq i} (\ln|\cos 2\pi a - \cos 2\pi\theta_j| - \ln|\cos 2\pi\theta_i - \cos 2\pi\theta_j|) \triangleq \sum_1 - \sum_2.$$

Lemma 2.8 reduces this problem to estimating the minimal terms.

First we estimate $\sum_1$:

$$\begin{aligned}\sum_1 &= \sum_{j\in I_1\cup I_2, j\neq i} \ln|\cos 2\pi a - \cos 2\pi\theta_j| \\ &= \sum_{j\in I_1\cup I_2, j\neq i} \ln|\sin\pi(a+\theta_j)| + \sum_{j\in I_1\cup I_2, j\neq i} \ln|\sin\pi(a-\theta_j)| + (6sq_n - 1)\ln 2 \\ &\triangleq \sum_{1,+} + \sum_{1,-} + (6sq_n - 1)\ln 2.\end{aligned}$$

We cut $\sum_{1,+}$ or $\sum_{1,-}$ into $6s$ sums and then apply Lemma 2.8, we get that for some absolute constant C_1:

$$\sum_1 \leq -6sq_n \ln 2 + C_1 s \ln q_n.$$

Next, we estimate $\sum_2$.

$$\begin{aligned}\sum_2 &= \sum_{j\in I_1\cup I_2, j\neq i} \ln|\cos 2\pi\theta_j - \cos 2\pi\theta_i| \\ &= \sum_{j\in I_1\cup I_2, j\neq i} \ln|\sin\pi(2\theta + (i+j)\alpha)| + \sum_{j\in I_1\cup I_2, j\neq i} \ln|\sin\pi(i-j)\alpha| + (6sq_n - 1)\ln 2 \\ &\triangleq \sum_{2,+} + \sum_{2,-} + (6sq_n - 1)\ln 2.\end{aligned}$$

We need to carefully estimate the minimal terms. For $\sum_{2,+}$, we use the property of resonant set; and for $\sum_{2,-}$, we use the Diophantine condition on α.

For any $0 < |j| < q_{n+1}$, we have $\|j\alpha\| \geq \|q_n\alpha\| \geq C_\xi e^{-\xi q_n}$. Therefore

$$\max(\ln|\sin x|, \ln|\sin(x + \pi j\alpha)|) \geq -2\xi q_n \ \text{ for } y > y(\alpha, \xi).$$

This means in any interval of length sq_n, there can be at most one term which is less than $-2\xi q_n$. Then there can be at most 6 such terms in total.

For the part $\sum_{2,-}$, since $\|(i-j)\alpha\| \geq C_\xi e^{-\xi|i-j|} \geq e^{-20\xi sq_n}$, these 6 smallest terms must be bounded by $-20\xi sq_n$ from below. Hence $\sum_{2,-} \geq -6sq_n\ln 2 - C\xi sq_n - Cs\ln q_n$ for $y > y(\xi)$ and some absolute constant C.

For the part $\sum_{2,+}$, notice $|i+j| \leq 2y + 4sq_n < 3y < |n_{j+1}|$ and $i + j > 0 > -n_j$. Suppose $\|2\theta + k_0\alpha\| = \min_{j\in I_1\cup I_2}\|2\theta + (i+j)\alpha\| \leq e^{-100\epsilon_0 sq_n} < e^{-\epsilon_0|k_0|}$. Then for any $|k| \leq |k_0| \leq 40sq_n$ (including $|n_j|$),

$$\|2\theta - k\alpha\| \geq \|(k+k_0)\alpha\| - \|2\theta + k_0\alpha\| > \|2\theta + k_0\alpha\| \ \text{ for } y > y(\alpha, \epsilon_0, \xi).$$

This means $-k_0$ must be a ϵ_0-resonance, therefore $|k_0| \leq |n_{j-1}|$. Then

$$\|2\theta - n_j\alpha\| \geq \|(n_j + k_0)\alpha\| - \|2\theta + k_0\alpha\| \geq C_\xi e^{-12\xi sq_n} - e^{-100\epsilon_0 sq_n} > e^{-100\epsilon_0 sq_n} \geq \|2\theta + k_0\alpha\|$$

leads to a contradiction. Thus the smallest terms must be greater than $-100\epsilon_0 sq_n$. We can bound $\sum_{2,+}$ by $-6sq_n\ln 2 - 600\epsilon_0 sq_n - 12\xi sq_n - Cs\ln q_n$ from below. Therefore $\sum_2 \geq -6sq_n\ln 2 -$

$C\epsilon_0 sq_n - C\xi sq_n - Cs\ln q_n$. Thus the set $\{\theta_j\}_{j\in I_1\cup I_2}$ is $C_4\epsilon_0 + C_4\xi-$uniform for $y > y(\alpha,\epsilon_0,\xi)$ and some absolute constant C_4. □

Now let C_4 be the absolute constant in Lemma 4.3. Choose $0 < 1000\xi < \epsilon_0 < \frac{\epsilon_1}{100C_4}$. Combining Lemma 4.2 and Lemma 4.3, we know that when $y > y(\alpha,\epsilon_0,\xi)$, $\{\theta_j\}_{j\in I_1\cup I_2}$ can not be inside the set $M_{6sq_n-1,\tilde{L}-2C_4\epsilon_0}$ at the same time. Therefore 0 and y can not be $(6sq_n-1,\tilde{L}-\int\ln|c_{\hat{\lambda}}|-9C_4\epsilon_0)$ at the same time. However 0 is $(6sq_n-1,\tilde{L}-\int\ln|c_{\hat{\lambda}}|-9C_4\epsilon_0)-$singular given n large enough. Therefore

$$\{\theta_j\}_{j\in I_1}\subset M_{6sq_n-1,\tilde{L}-2C_4\epsilon_0}.$$

Thus y must be $(6sq_n-1,\tilde{L}-\int\ln|c_{\hat{\lambda}}|-9C_4\epsilon_0)-$regular. This implies

$$|u(y)|\le e^{-(\tilde{L}-\int\ln|c_{\hat{\lambda}}|-9C_4\epsilon_0)\frac{1}{4}|y|} < e^{-\frac{\epsilon_1}{5}|y|} \text{ for } |y|\ge y(\lambda,\alpha,\epsilon_0,\xi).$$

Thus there exists $C_3 = C_{\lambda,\alpha,\epsilon_0,\xi}$ such that $|u(y)|\le C_3e^{-\frac{\epsilon_1}{5}|y|}$ for any $3|n_j|\le|y|\le\frac{1}{3}|n_{j+1}|$ and $j\in\mathbb{N}$.

5. Almost reducibility in region II

Proof of Theorem 3.3

For any $E\in\Sigma_\lambda$, take $\theta(E)$ and $\{u_k\}$ as in Theorem 2.6. Let ϵ_1 be as in (2.4), C_4 be the absolute constant from Lemma 4.3, and C_2 be the absolute constant from Lemma 2.9. Fix $\max(32C_2\xi,1000\xi) < \epsilon_0 < \min(\frac{\epsilon_1}{200},\frac{\epsilon_1}{100C_4})$. By Lemma 3.2, there exists C depending on λ and α such that for any $3|n_j|<|k|<\frac{1}{3}|n_{j+1}|$, we have $|u_k|\le Ce^{-\frac{\epsilon_1}{5}|k|}$.

For any n, $9|n_j|<n<\frac{1}{9}|n_{j+1}|$, of the form

$$n = rq_m - 1 < q_{m+1}.\ ^{2} \tag{5.1}$$

Let $u(x)=u^I(x)=\sum_{k\in I}u_ke^{2\pi ikx}$ with $I=[-[\frac{n}{2}],[\frac{n}{2}]]=[x_1,x_2]$. Define

$$U(x)=\begin{pmatrix}e^{2\pi i\theta}u(x)\\ u(x-\alpha)\end{pmatrix}.$$

Let $A(\theta)=A_{\lambda,E}(\theta)$. By direct computation:

$$A(x)U(x)=e^{2\pi i\theta}U(x+\alpha)+\begin{pmatrix}g(x)\\0\end{pmatrix}\triangleq e^{2\pi i\theta}U(x+\alpha)+G(x).$$

The Fourier coefficients of $g(x)$ are possibly nonzero only at four points x_1, x_2, x_1-1 and x_2+1. Since $|u_k|\le C_1e^{-\frac{\epsilon_1}{5}|k|}$ when $3|n_j|<|k|<\frac{1}{3}|n_{j+1}|$, we know that $\|G(x)\|_{\frac{\epsilon_1}{20\pi}}\le C_1e^{-\frac{\epsilon_1}{20}n}$.

Combining Lemma $A.3$ and 2.1, we have exponential control of the growth of the transfer matrix, for any $\delta>0$ there exists $C_\delta>0$ such that

$$\|\tilde{A}_k(x)\|_{\frac{\epsilon_1}{2\pi}}\le C_\delta e^{\delta|k|},\quad\text{for any } k.$$

With some effort we are able to get the following significantly improved upper bound:

Theorem 5.1. *For some $C>0$ depending on λ and α,*

$$\|\tilde{A}_k(x)\|_{\mathbb{T}}\le C(1+|k|)^C.$$

[2]The existence of such n comes from (4.1).

Proof.

Let $\tilde{U}(x) = Q(x)U(x)$, $\tilde{G}(x) = Q(x+\alpha)G(x)$, where $Q = Q_\lambda$ is given in $(A.2)$. Since

$$\max\left(\|Q(x)\|_{\frac{\epsilon_1}{20\pi}}, \|Q^{-1}(x)\|_{\frac{\epsilon_1}{20\pi}}\right) \leq C,$$

we have

$$\tilde{A}(x)\tilde{U}(x) = e^{2\pi i\theta}\tilde{U}(x+\alpha) + \tilde{G}(x),$$

where $\|\tilde{G}(x)\|_{\frac{\epsilon_1}{20\pi}} \leq Ce^{-\frac{\epsilon_1}{20}n}$.

Lemma 5.2. *Let C_2 be the constant from Lemma 2.9, then for any δ, $2C_2\xi < \delta < \frac{\epsilon_0}{16}$, we have*

$$\inf_{|\mathrm{Im}(x)|\leq\frac{\epsilon_1}{20\pi}} \|\tilde{U}(x)\| \geq e^{-2\delta n},$$

for $n > n(\alpha, \delta)$.

Proof. We will prove the statement by contradiction. Suppose for some $x_0 \in \{|\mathrm{Im}(x)| \leq \frac{\epsilon_1}{20\pi}\}$ we have $\|\tilde{U}(x_0)\| < e^{-2\delta n}$. Notice that for any $l \in \mathbb{N}$,

$$e^{2\pi il\theta}\tilde{U}(x_0+l\alpha) = \tilde{A}_l(x_0)\tilde{U}(x_0) - \sum_{m=1}^{l} e^{2\pi i(m-1)\theta}\tilde{A}_{l-m}(x_0+m\alpha)\tilde{G}(x_0+(m-1)\alpha).$$

This implies for $n > n(\delta)$ large enough and for any $0 \leq l \leq n$, $\|\tilde{U}(x_0+l\alpha)\| \leq e^{-\delta n}$, thus $\|u(x_0+l\alpha)\| \leq C_\delta e^{-\delta n}$. By Lemma 2.9, $\|u(x+i\mathrm{Im}(x_0))\|_{\mathbb{T}} \leq C_2C_\delta e^{C_2\xi n}e^{-\delta n} \leq e^{-\frac{\delta}{2}n}$. This contradicts with $\int_{\mathbb{T}} u(x+i\mathrm{Im}(x_0))\mathrm{d}x = u_0 = 1$. □

Lemma 5.3. [3] *Let $V : \mathbb{T} \to \mathbb{C}^2$ be analytic in $|\mathrm{Im}(x)| < \eta$. Assume that $\delta_1 < \|V(x)\| < \delta_2^{-1}$ holds on $|\mathrm{Im}(x)| < \eta$. Then there exists $M : \mathbb{T} \to SL(2,\mathbb{C})$ analytic on $|\mathrm{Im}(x)| < \eta$ with first column V and $\|M\|_\eta \leq C\delta_1^{-2}\delta_2^{-1}(1-\ln(\delta_1\delta_2))$.*

Applying Lemma 5.3, let $M(x)$ be the matrix with first column $\tilde{U}(x)$. Then $e^{-2\delta n} \leq \|\tilde{U}(x)\|_{\frac{\delta}{\pi}} \leq e^{\delta n}$ and hence $\|M(x)\|_{\frac{\delta}{\pi}} \leq Ce^{6\delta n}$. Therefore

$$M^{-1}(x+\alpha)\tilde{A}(x)M(x) = \begin{pmatrix} e^{2\pi i\theta} & 0 \\ 0 & e^{-2\pi i\theta} \end{pmatrix} + \begin{pmatrix} \beta_1(x) & b(x) \\ \beta_3(x) & \beta_4(x) \end{pmatrix}$$

where $\|\beta_1(x)\|_{\frac{\delta}{\pi}}$, $\|\beta_3(x)\|_{\frac{\delta}{\pi}}$, $\|\beta_4(x)\|_{\frac{\delta}{\pi}} \leq Ce^{-\frac{\epsilon_1}{40}n}$, and $\|b(x)\|_{\frac{\delta}{\pi}} \leq Ce^{13\delta n}$. Let

$$\Phi(x) = M(x)\begin{pmatrix} e^{\frac{\epsilon_1}{160}n} & 0 \\ 0 & e^{-\frac{\epsilon_1}{160}n} \end{pmatrix}.$$

Then we would have:

$$\Phi(x+\alpha)^{-1}\tilde{A}(x)\Phi(x) = \begin{pmatrix} e^{2\pi i\theta} & 0 \\ 0 & e^{-2\pi i\theta} \end{pmatrix} + H(x),$$

where $\|H(x)\|_{\frac{\delta}{\pi}} \leq Ce^{-\frac{\epsilon_1}{160}n}$, and $\|\Phi(x)\|_{\frac{\delta}{\pi}} \leq Ce^{\frac{\epsilon_1}{80}n}$. Thus

$$\sup_{0\leq s\leq e^{\frac{\epsilon_1}{320}n}} \|\tilde{A}_s(x)\|_{\mathbb{T}} \leq e^{\frac{\epsilon_1}{20}n}$$

for $n \geq n(\lambda,\alpha)$ satisfying (5.1). For s large, there always exists $9|n_j| < n < \frac{1}{9}|n_{j+1}|$ satisfying (5.1) such that $cn \leq \frac{320}{\epsilon_1}\ln s \leq n$ with some absolute constant c. Thus there exists C depending on λ and α such that $\|\tilde{A}_k(x)\|_{\mathbb{T}} \leq C(1+|k|)^C$. □

Now we come back to the proof of Theorem 3.3. Fix some $n = |n_j|$, and $N = |n_{j+1}|$. Let $u(x) = u^{I_2}(x)$ with $I_2 = [-[\frac{N}{9}],[\frac{N}{9}]]$ and $U(x) = \begin{pmatrix} e^{2\pi i\theta}u(x) \\ u(x-\alpha)\end{pmatrix}$. Then

$$A(x)U(x) = e^{2\pi i\theta}U(x+\alpha) + G(x) \quad with \quad \|G(x)\|_{\frac{\epsilon_1}{20\pi}} \leq Ce^{-\frac{\epsilon_1}{90}N}.$$

Define $U_0(x) = e^{\pi i n_j x}U(x)$. Notice that if n_j is even, then $U_0(x)$ is well-defined on $\mathbb{T}$, otherwise $U_0(x+1) = -U_0(x)$.

$$\tilde{A}(x)\tilde{U}_0(x) = e^{2\pi i\tilde{\theta}}\tilde{U}_0(x+\alpha) + H(x),$$

where $\tilde{\theta} = \theta - \frac{n_j}{2}\alpha$, $\tilde{U}_0(x) = Q(x)U_0(x)$ and $\|H(x)\|_{\frac{\epsilon_1}{20\pi}} \leq Ce^{-\frac{\epsilon_1}{100}N}$. Consider the matrix $W(x)$ with $\tilde{U}_0(x)$ and $\overline{\tilde{U}_0(x)}$ being its two columns. Then

$$\tilde{A}(x)W(x) = W(x+\alpha)\begin{pmatrix} e^{2\pi i\tilde{\theta}} & 0 \\ 0 & e^{-2\pi i\tilde{\theta}}\end{pmatrix} + \tilde{H}(x).$$

Theorem 5.4. *Let $L^{-1} = \|2\theta - n_j\alpha\|$. Then for $n > N_0(\lambda,\alpha)$ we have*

$$|\det W(x)| \geq L^{-4C} \quad \text{for any } x \in \mathbb{T},$$

where C is the constant appeared in Theorem 5.1.

Proof. First, we fix $\xi_1 < \frac{\epsilon_0}{1600}$ so that $\|k\alpha\| \geq C_{\xi_1}e^{-\xi_1|k|}$ for any $k \neq 0$. We have the following estimate about L:

Lemma 5.5. $e^{\epsilon_0 n} \leq L \leq e^{4\xi_1 N}$.

$$e^{-2\xi_1 N} \leq \|(n_{j+1}-n_j)\alpha\| \leq 2\|n_j\alpha - 2\theta\| = 2L^{-1} \leq 2e^{-\epsilon_0 n} \quad \text{for } n \geq N(\xi_1).$$

Now we prove by contradiction. Suppose there exists κ and $x_0 \in \mathbb{T}$ such that $\|\tilde{U}_0(x_0) - \kappa\overline{\tilde{U}_0(x_0)}\| < L^{-4C}$. Then

$$\begin{aligned}
&\|\tilde{U}_0(x_0+l\alpha)e^{2\pi il\tilde{\theta}} - \kappa\overline{\tilde{U}_0(x_0+l\alpha)}e^{-2\pi il\tilde{\theta}}\| \\
\leq&\|\sum_{m=0}^{l-1}\tilde{A}_{l-m}(x_0+m\alpha)H(x_0+m\alpha) - \kappa\sum_{m=0}^{l-1}\tilde{A}_{l-m}(x_0+m\alpha)\overline{H(x_0+m\alpha)}\| + \|A_l(x_0)\|L^{-4C} \\
\leq&CL^{2C}e^{-\frac{\epsilon_1}{100}N} + CL^{-2C} < L^{-C}.
\end{aligned}$$

for $0 \leq |l| \leq L^2$. If we take $j = \frac{L}{4}$, then

$$\|\tilde{U}_0(x_0+\frac{L}{4}\alpha) + \kappa\overline{\tilde{U}_0(x_0+\frac{L}{4}\alpha)}\| < L^{-1}. \tag{5.2}$$

Next since $\|U_0(x)\|_{\mathbb{T}} \leq n$, we have $\|\tilde{U}_0(x)\|_{\mathbb{T}} \leq Cn$. Thus

$$\|\tilde{U}_0(x_0+l\alpha) - \kappa\overline{\tilde{U}_0(x_0+l\alpha)}\| < L^{-\frac{1}{3}} \quad \text{for } 0 \leq |l| \leq L^{\frac{1}{2}}.$$

For any analytic function $f(x)=\sum_{k\in\mathbb{Z}}\hat{f}_k e^{2\pi ikx}$, define $f_{[-m,m]}(x)=\sum_{|k|\le m}\hat{f}_k e^{2\pi ikx}$. For any column vector $V(x)=\begin{pmatrix}v^{(1)}(x)\\ v^{(2)}(x)\end{pmatrix}$, let $V_{[-m,m]}(x)=\begin{pmatrix}v^{(1)}_{[-m,m]}(x)\\ v^{(2)}_{[-m,m]}(x)\end{pmatrix}$. Now let us define $\tilde{U}_0^{[9n]}(x)=Q(x)e^{\pi in_j x}U_{[-9n,9n]}(x)$. Then

$$\|\tilde{U}_0^{[9n]}(x)-\tilde{U}_0(x)\|_{\mathbb{T}}\le Ce^{-\frac{9}{5}\epsilon_1 n}.$$

Consider $[e^{-\pi in_j x}\tilde{U}_0^{[9n]}(x)]_{[-18n,18n]}(x)e^{\pi in_j x}$. This function differs from a polynomial with essential degree $36n$ only by a multiple of $e^{\pi in_j x}$. Notice that $Q(x)$ is analytic in $\{x:|\mathrm{Im}(x)|\le\frac{\epsilon_1}{4\pi}\}$, thus $|\hat{Q}(k)|\le Ce^{-\frac{\epsilon_1}{2}|k|}$. Then

$$|\widehat{e^{-\pi in_j x}\tilde{U}_0^{[9n]}}(k)|\le\sum_{|m|\le 9n}|\hat{Q}(k-m)\hat{U}(m)|\le Cne^{-\frac{\epsilon_1}{2}(|k|-9n)}\ \text{ for } |k|\ge 18n.$$

Thus

$$\|e^{-\pi in_j x}\tilde{U}_0^{[9n]}(x)-[e^{-\pi in_j x}\tilde{U}_0^{[9n]}]_{[-18n,18n]}(x)\|_{\mathbb{T}}\le e^{-4\epsilon_1 n},$$

$$\|\tilde{U}_0(x)-[e^{-\pi in_j x}\tilde{U}_0^{[9n]}]_{[-18n,18n]}(x)e^{\pi in_j x}\|_{\mathbb{T}}\le e^{-4\epsilon_1 n}.$$

Hence

$$\begin{aligned}&\|[e^{-\pi in_j x}\tilde{U}_0^{[9n]}]_{[-18n,18n]}(x_0+l\alpha)e^{2\pi in_j(x_0+l\alpha)}-\kappa\overline{[e^{-\pi in_j x}\tilde{U}_0^{[9n]}]_{[-18n,18n]}(x_0+l\alpha)}\|_{\mathbb{T}}\\ <&2L^{-\frac{1}{3}}+e^{-4\epsilon_1 n},\end{aligned}$$

for $|l|\le L^{\frac{1}{2}}$. Notice that

$$[e^{-\pi in_j x}\tilde{U}_0^{[9n]}]_{[-18n,18n]}(x)e^{2\pi in_j x}-\kappa\overline{[e^{-\pi in_j x}\tilde{U}_0^{[9n]}]_{[-18n,18n]}(x)}$$

is a polynomial whose essential degree is at most $37n$. Thus by Lemma 2.9, we would have

$$\|[e^{-\pi in_j x}\tilde{U}_0^{[9n]}]_{[-18n,18n]}(x)e^{\pi in_j x}-\kappa\overline{[e^{-\pi in_j x}\tilde{U}_0^{[9n]}]_{[-18n,18n]}(x)e^{\pi in_j x}}\|_{\mathbb{T}}<L^{-\frac{1}{4}}+e^{-2\epsilon_1 n}.$$

Hence $\|\tilde{U}_0(x)-\kappa\overline{\tilde{U}_0(x)}\|_{\mathbb{T}}<L^{-\frac{1}{4}}+2e^{-2\epsilon_1 n}$. But combining with (9.1) we would get $\|\tilde{U}_0(x_0+\frac{L}{4}\alpha)\|<2L^{-\frac{1}{4}}+2e^{-2\epsilon_1 n}$, but this contradicts with $\inf_{x\in\mathbb{T}}\|\tilde{U}_0(x)\|>e^{-2\delta n}$ since $\delta<\frac{\epsilon_0}{16}$. □

Now for $n>N_0(\lambda,\alpha)$, take $S(x)=\mathrm{Re}\tilde{U}_0(x)$ and $T(x)=\mathrm{Im}\tilde{U}_0(x)$. Let $W_1(x)$ be the matrix with columns $S(x)$ and $T(x)$. Notice that $\det W_1(x)$ is well-defined on $\mathbb{T}$ and $\det W_1(x)\neq 0$ on $\mathbb{T}$, hence without loss of generality we could assume $\det W_1(x)>0$ on $\mathbb{T}$, otherwise we simply take $W_1(x)$ to be the matrix with columns $S(x)$ and $-T(x)$. Then

$$\|\tilde{A}(x)W_1(x)-W_1(x+\alpha)R_{-\tilde{\theta}}\|_{\mathbb{T}}\le Ce^{-\frac{\epsilon_1}{45}N}.$$

By taking determinant, we get

$$\det W_1(x)=\det W_1(x+\alpha)+O(e^{-\frac{\epsilon_1}{50}N})\ \text{ on } \mathbb{T}.$$

Since $\det W_1(x)$ is analytic on $|\mathrm{Im}x|\le\frac{\epsilon_1}{20\pi}$, by considering the Fourier coefficients we could get

$$\det W_1(x)=w_0+O(e^{-\frac{\epsilon_1}{100}N})\ \text{ on } \mathbb{T},$$

where $w_0\ge L^{-5C}$. Thus $\det W_1(x)$ is almost a positive constant.

Define $W_2(x)=\det W_1(x)^{-\frac{1}{2}}W_1(x)$. Then $W_2(x)\in C^\omega(\mathbb{T})$ and $\det W_2(x)=1$. We have

$$W_2^{-1}(x+\alpha)\tilde{A}(x)W_2(x)=\frac{\det W_1(x+\alpha)^{\frac{1}{2}}}{\det W_1(x)^{\frac{1}{2}}}R_{-\tilde{\theta}}+O(e^{-\frac{\epsilon_1}{100}N})\ \text{ on } \mathbb{T},$$

$$W_2^{-1}(x+\alpha)\tilde{A}(x)W_2(x) = R_{-\tilde{\theta}} + O(e^{-\frac{\epsilon_1}{200}N}) \text{ on } \mathbb{T}.$$

Now let's prove $\deg W_2(x) \leq 36n$. $\deg W_2(x)$ is the same as the degree of its columns. For $M : \mathbb{R}/2\mathbb{Z} \to \mathbb{R}^2$, we say $\deg M = k$ if M is homotopic to $\begin{pmatrix}\cos k\pi x\\ \sin k\pi x\end{pmatrix}$.

For some constant $c > 0$, we obviously have

$$\int_{\mathbb{T}} \|S(x)\| \, \mathrm{d}x + \int_{\mathbb{T}} \|T(x)\| \, \mathrm{d}x \geq \int_{\mathbb{T}} \|S(x) + iT(x)\| \, \mathrm{d}x = \int_{\mathbb{T}} \|\tilde{U}_0(x)\| \, \mathrm{d}x \geq c.$$

Without loss of generality we could assume $\int_{\mathbb{T}} \|S(x)\| \, \mathrm{d}x > \frac{c}{2}$. Also

$$\tilde{A}(x)S(x) = S(x+\alpha)\cos 2\pi\tilde{\theta} - T(x+\alpha)\sin 2\pi\tilde{\theta} + O(e^{-\frac{\epsilon_1}{45}N}) \text{ on } \mathbb{T}.$$

Then since $\|2\tilde{\theta}\| = L^{-1}$,

$$\tilde{A}(x)S(x) = S(x+\alpha) + O(L^{-\frac{1}{2}}) \text{ on } \mathbb{T}.$$

First we prove $\inf_{x\in\mathbb{T}} \|S(x)\| \geq e^{-2\epsilon_1 n}$. Suppose otherwise. Then there exists $x_0 \in \mathbb{T}$, so that $\|S(x_0)\| < e^{-2\epsilon_1 n}$. Then $\|\mathrm{Re}\tilde{U}_0(x_0 + l\alpha)\| < e^{-\frac{\epsilon_0}{8}n}$ for $|l| < e^{\frac{\epsilon_0}{4C}n}$, where C is the constant that appeared in Theorem 5.1. We have already shown that

$$\|\tilde{U}_0(x) - [e^{-\pi i n_j x}\tilde{U}_0^{[9n]}]_{[-18n,18n]}e^{\pi i n_j x}\|_{\mathbb{T}} < e^{-4\epsilon_1 n}.$$

Thus

$$\|\mathrm{Re}[e^{-\pi i n_j x}\tilde{U}_0^{[9n]}]_{[-18n,18n]}(x_0 + l\alpha)\| < e^{-\frac{\epsilon_0}{16}n}$$

for $|l| < e^{\frac{\epsilon_0}{4C}n}$. However $\mathrm{Re}[e^{-\pi i n_j x}\tilde{U}_0^{[9n]}]_{[-18n,18n]}$ is a polynomial with essential degree at most $36n$. Using Lemma 2.9 we are able to get $\|\mathrm{Re}[e^{-\pi i n x}\tilde{U}_0^{[9n]}]_{[-18n,18n]}e^{\pi i n_j x}\|_{\mathbb{T}} < e^{-\frac{\epsilon_0}{32}n}$, and thus $\|\mathrm{Re}\tilde{U}_0(x)\|_{\mathbb{T}} < e^{-\frac{\epsilon_0}{64}n}$ which is a contradiction to $\int_{\mathbb{T}} \|\mathrm{Re}\tilde{U}_0(x)\| \, \mathrm{d}x > \frac{c}{2}$. At the meantime, we also get $\|S(x) - \mathrm{Re}[e^{-\pi i n_j x}\tilde{U}_0^{[9n]}]_{[-18n,18n]}(x)e^{\pi i n_j x}\|_{\mathbb{T}} \triangleq \|S(x) - h(x)\|_{\mathbb{T}} \leq e^{-4\epsilon_1 n}$. The first column of $W_2(x)$ is $\det W_1(x)^{-\frac{1}{2}}S(x)$. We have

$$\begin{aligned}
&\|\frac{S(x)}{\det W_1(x)^{\frac{1}{2}}} - \frac{h(x)}{{w_0}^{\frac{1}{2}}}\| \\
\leq &\frac{1}{|\det W_1(x)^{\frac{1}{2}}|}\|S(x) - h(x) + (1 - \frac{\det W_1(x)^{\frac{1}{2}}}{{w_0}^{\frac{1}{2}}})h(x)\| \\
\leq &L^{2C}(e^{-4\epsilon_1 n} + L^{8C}e^{-\frac{\epsilon_1}{100}N}) \\
\leq &e^{-3\epsilon_1 n} < \|\frac{S(x)}{\det W_1(x)^{\frac{1}{2}}}\| \text{ on } \mathbb{T}.
\end{aligned}$$

Thus by Rouché's theorem $|\deg W_2(x)| = |\deg h(x)| \leq 19n$. Notice that

$$|\rho(\alpha, W_2^{-1}\tilde{A}W_2) + \tilde{\theta}| < Ce^{-\frac{\epsilon_1}{200}N}.$$

Then, by 2.2 for some $|m| \leq 19n$:

$$|\rho(\alpha, \tilde{A}) - \frac{m}{2}\alpha + \tilde{\theta}| < Ce^{-\frac{\epsilon_1}{200}N}.$$

Appendix A.

When λ belongs to region II, let $\epsilon_2 = \ln \frac{\lambda_2+\sqrt{\lambda_2^2-4\lambda_1\lambda_3}}{\lambda_1+\lambda_3+\sqrt{(\lambda_1+\lambda_3)^2-4\lambda_1\lambda_3}} > \epsilon_1$. Then $c(x)$ is analytic and nonzero on $|\mathrm{Im}(x)| < \frac{\epsilon_2}{2\pi}$. Furthermore, the winding number of $c(\cdot + i\epsilon)$ is equal to zero when $|\epsilon| < \frac{\epsilon_2}{2\pi}$.

Lemma A.1. *When λ belongs to region II, we can find an analytic function $f(x)$ on $|\mathrm{Im}(x)| \le \frac{\epsilon_1}{2\pi}$ such that $c(x) = |c|(x)e^{f(x+\alpha)-f(x)}$ and $\tilde{c}(x) = |c|(x)e^{-f(x+\alpha)+f(x)}$.*

Proof. Since the winding numbers of $c(x)$ and $\tilde{c}(x)$ are 0 on $|\mathrm{Im}(x)| \le \frac{\epsilon_1}{2\pi}$, there exist analytic functions $g_1(x)$ and $g_2(x)$ on $|\mathrm{Im}(x)| \le \frac{\epsilon_1}{2\pi}$, such that $c(x) = e^{g_1(x)}$ and $\tilde{c}(x) = e^{g_2(x)}$. Notice that

$$\int_{\mathbb{T}} \ln|c(x)|\,\mathrm{d}x = \int_{\mathbb{T}} \ln|\tilde{c}(x)|\,\mathrm{d}x$$
$$\int_{\mathbb{T}} \arg c(x)\,\mathrm{d}x = \int_{\mathbb{T}} \arg \tilde{c}(x)\,\mathrm{d}x,$$

so there exists an analytic function $f(x)$ such that $2f(x+\alpha) - 2f(x) = g_1(x) - g_2(x)$. Then $c(x) = |c|(x)e^{f(x+\alpha)-f(x)}$. □

Lemma A.2. *When λ belongs to region II, there exists an analytic matrix $Q_\lambda(x)$ defined on $|\mathrm{Im}(x)| \le \frac{\epsilon_1}{2\pi}$ such that*

$$Q_\lambda^{-1}(x+\alpha)\tilde{A}_{\lambda,E}(x)Q_\lambda(x) = A_{\lambda,E}(x).$$

Proof.

$$\begin{aligned}
\tilde{A}_{\lambda,E}(x) =& \frac{1}{\sqrt{|c|(x)|c|(x-\alpha)}} \begin{pmatrix} 1 & 0 \\ 0 & \sqrt{\frac{\tilde{c}(x)}{c(x)}} \end{pmatrix} \begin{pmatrix} E-v(x) & -\tilde{c}(x-\alpha) \\ c(x) & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & \sqrt{\frac{c(x-\alpha)}{\tilde{c}(x-\alpha)}} \end{pmatrix} \\
=& \frac{c(x)}{\sqrt{|c|(x)|c|(x-\alpha)}} \begin{pmatrix} 1 & 0 \\ 0 & \sqrt{\frac{\tilde{c}(x)}{c(x)}} \end{pmatrix} A(x) \begin{pmatrix} 1 & 0 \\ 0 & \sqrt{\frac{c(x-\alpha)}{\tilde{c}(x-\alpha)}} \end{pmatrix} \\
=& e^{f(x+\alpha)}\sqrt{|c|(x)} \begin{pmatrix} 1 & 0 \\ 0 & \sqrt{\frac{\tilde{c}(x)}{c(x)}} \end{pmatrix} A(x) \left\{ e^{f(x)}\sqrt{|c|(x-\alpha)} \begin{pmatrix} 1 & 0 \\ 0 & \sqrt{\frac{\tilde{c}(x-\alpha)}{c(x-\alpha)}} \end{pmatrix} \right\}^{-1} \\
=& Q_\lambda(x+\alpha)A_{\lambda,E}(x)Q_\lambda^{-1}(x).
\end{aligned}$$

□

Lemma A.3. *If α is irrational, λ belongs to region II, $E \in \Sigma(\lambda)$, then $L(\alpha, A_{\lambda,E}(\cdot + i\epsilon)) = L(\alpha, \tilde{A}_{\lambda,E}(\cdot + i\epsilon)) = 0$ for $|\epsilon| \le \frac{\epsilon_1}{2\pi}$.*

Proof. $L(A(\cdot + i\epsilon)) = L(D(\cdot + i\epsilon)) - \int \ln|c(x+i\epsilon)|\mathrm{d}x$

$$\begin{aligned}
D(x+i\epsilon) &= \begin{pmatrix} E - e^{2\pi i(x+i\epsilon)} - e^{-2\pi i(x+i\epsilon)} & -\lambda_1 e^{2\pi i(x-\frac{\alpha}{2}+i\epsilon)} - \lambda_2 - \lambda_3 e^{-2\pi i(x-\frac{\alpha}{2}+i\epsilon)} \\ \lambda_1 e^{-2\pi i(x+\frac{\alpha}{2}+i\epsilon)} + \lambda_2 + \lambda_3 e^{2\pi i(x+\frac{\alpha}{2}+i\epsilon)} & 0 \end{pmatrix} \\
&= e^{2\pi\epsilon} \begin{pmatrix} -e^{2\pi ix} + o(1) & -\lambda_3 e^{-2\pi i(x-\frac{\alpha}{2})} + o(1) \\ \lambda_1 e^{-2\pi i(x+\frac{\alpha}{2})} + o(1) & 0 \end{pmatrix}.
\end{aligned}$$

Thus the asymptotic behaviour of $L(D(\cdot+i\epsilon))$ is:

$$L(D(\cdot+i\epsilon))=\ln|\frac{1+\sqrt{1-4\lambda_1\lambda_3}}{2}|+2\pi\epsilon \quad \text{when } \epsilon\to\infty,$$
$$L(D(\cdot+i\epsilon))=\ln|\frac{1+\sqrt{1-4\lambda_1\lambda_3}}{2}|-2\pi\epsilon \quad \text{when } \epsilon\to-\infty.$$

Then it suffices to calculate $\int\ln|c(x+i\epsilon)|\mathrm{d}x$ in region II. We have

$$\begin{aligned}&\int\ln|c(x+i\epsilon)|\mathrm{d}x\\ &=\ln\lambda_3-2\pi\epsilon+\int\ln|e^{2\pi ix}-y_{1,\epsilon}|+\int\ln|e^{2\pi ix}-y_{2,\epsilon}|.\end{aligned}$$

where $y_{1,\epsilon}=\frac{-\lambda_2+\sqrt{\lambda_2^2-4\lambda_1\lambda_3}}{2\lambda_3}e^{2\pi\epsilon}$ and $y_{2,\epsilon}=\frac{-\lambda_2-\sqrt{\lambda_2^2-4\lambda_1\lambda_3}}{2\lambda_3}e^{2\pi\epsilon}$.

$$\int\ln|c(x+i\epsilon)|\mathrm{d}x=\begin{cases}2\pi\epsilon+\ln\lambda_1 & \epsilon>\frac{1}{2\pi}\ln\frac{\lambda_2+\sqrt{\lambda_2^2-4\lambda_1\lambda_3}}{2\lambda_1},\\ \ln\frac{\lambda_2+\sqrt{\lambda_2^2-4\lambda_1\lambda_3}}{2} & \frac{1}{2\pi}\ln\frac{\lambda_2-\sqrt{\lambda_2^2-4\lambda_1\lambda_3}}{2\lambda_1}\le\epsilon\le\frac{1}{2\pi}\ln\frac{\lambda_2+\sqrt{\lambda_2^2-4\lambda_1\lambda_3}}{2\lambda_1},\\ -2\pi\epsilon+\ln\lambda_3 & \epsilon<\frac{1}{2\pi}\ln\frac{\lambda_2-\sqrt{\lambda_2^2-4\lambda_1\lambda_3}}{2\lambda_1}.\end{cases}$$

Thus $L(A(\cdot+i\epsilon))=0$ when $|\epsilon|\le\frac{1}{2\pi}\ln\frac{\lambda_2+\sqrt{\lambda_2^2-4\lambda_1\lambda_3}}{\max(1,\lambda_1+\lambda_3)+\sqrt{\max(1,\lambda_1+\lambda_3)^2-4\lambda_1\lambda_3}}=\frac{\epsilon_1}{2\pi}$.

Since $\tilde{A}_{\lambda,E}(x+i\epsilon)=Q_\lambda(x+\alpha+i\epsilon)A_{\lambda,E}(x+i\epsilon)Q_\lambda^{-1}(x+i\epsilon)$, the statement about $\tilde{A}_{\lambda,E}$ is also true. □

Acknowledgement

I am deeply grateful to Svetlana Jitomirskaya for suggesting this problem and for many valuable discussions. This research was partially supported by the NSF DMS1401204.

References

1. S. Aubry and G. André, Analyticity breaking and Anderson localization in incommensurate lattices, In: Group Theoretical Methods in Physics (Kiryat Anavim, 1979), Ann. Israel Phys. Soc. 3, Hilger, Bristol, 133164 (1980) MR 83b:82076
2. A. Avila, Absolutely continuous spectrum for the almost Mathieu operator. Preprint
3. A. Avila, Almost reducibility and absolute continuity I. Preprint
4. A. Avila, Global theory of one-frequency Schrödinger operators, Acta Math. 215 (2015), no. 1, 154
5. A. Avila and S. Jitomirskaya, The Ten Martini Problem, Ann. of Math. (2) 170 (2009), no. 1, 303342, MR 2011a:47081
6. A. Avila and S. Jitomirskaya, Almost localization and almost reducibility, J. Eur. Math. Soc. 12 (2010), no. 1, 93131 MR 2011d:47071
7. A. Avila, S. Jitomirskaya and C. Marx, Spectral theory of extended Harper's model and a question by Erdős and Szekeres. Preprint
8. J. Avron and B. Simon, Almost periodic Schrödinger operators. II. The integrated density of states, Duke Math. J. 50 (1983), no. 1, 369391 MR 85i:34009a
9. A. Avila, J. You and Q. Zhou, Dry Ten Martini problem in non-critical case, in preparation
10. J. Bellissard, R. Lima and D. Testard, Almost periodic Schrdinger operators. Mathematics + physics. Vol. 1, 164, World Sci. Publishing, Singapore, 1985. MR 87m:46142
11. Y. Berezanskii, Expansions in eigenfunctions of selfadjoint operators. Transl. Math. Monogr., Vol. 17. Providence, RI: Am. Math. Soc. (1968)
12. M. D. Choi, G. A. Elliott and N. Yui, Gauss polynomials and the rotation algebra, Invent. Math. 99 (1990), no. 2, 225246 MR 91b:46067

13. S. Jitomirskaya, D.A. Koslover and M.S. Schulteis, Localization for a family of one-dimensional quasiperiodic operators of magnetic origin, Ann. Henri Poincar 6 (2005), no. 1, 103124, MR 2005j:81057
14. S. Jitomirskaya and C. Marx, Analytic quasi-perodic cocycles with singularities and the Lyapunov exponent of extended Harper's model, Comm. Math. Phys. 316 (2012), no. 1, 237267
15. R. Johnson and J. Moser, The rotation number for almost periodic potentials, Comm. Math. Phys. 84 (1982), no. 3, 403438 MR 83h:34018
16. J. Puig, Cantor spectrum for the almost Mathieu operator, Comm. Math. Phys. 244 (2004), 297-309 MR 2004k:11129
17. B. Simon, Schrödinger semigroups, Bull. Amer. Math. Soc. (N.S.) 7 (1982), no. 3, 447526 MR 86b:81001a

Department of Mathematics, University of California, Irvine, CA, 92697-3875, United States of America

E-mail address: rhan2@uci.edu